

AFCRL-68-0673

CLIMATIC NORMALS AS PREDICTORS Part 2: Extension

by

1 NOVEMBER 1968

ARNOLD COURT
PROFESSOR of CLIMATOLOGY
SAN FERNANDO VALLEY STATE COLLEGE



SCIENTIFIC REPORT NO 2

CONTRACT AF19 (628)-5176

PROJECT NO. 8624 TASK NO. 862402

WORK UNIT NO. 86240201

REPORT NO. 68-82-2

Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.

SAN FERNANDO VALLEY
STATE COLLEGE
FOUNDATION

NORTHRIDGE
(Los Angeles)
Calif. 91324

AD 686163

CONTRACT MONITOR: IVER A. LUND, CREW
Aerospace Instrumentation Laboratory

prepared for

AIR FORCE CAMBRIDGE RESEARCH LABORATORIES
OFFICE OF AEROSPACE RESEARCH
UNITED STATES AIR FORCE
BEDFORD, MASSACHUSETTS
01730

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va. 22151

AFCRL-68-0673

CLIMATIC NORMALS AS PREDICTORS

PART 2: EXTENSION

BY

ARNOLD COURT
Professor of Climatology
San Fernando Valley State College
Northridge (Los Angeles) Calif., 91324

Scientific Report No. 2

Contract AF 19(628)-5716

Project No. 8624 Task No. 862402
Work Unit No. 86240201

1 November 1968

Contract Monitor: Iver A. Lund
Aerospace Instrumentation Laboratory

Distribution of this document is unlimited. It may be released to the
Clearinghouse, Department of Commerce, for sale to the general public

Prepared for

Air Force Cambridge Research Laboratories

Office of Aerospace Research

United States Air Force

Bedford, Massachusetts

01730

ABSTRACT

Temperature and precipitation for a single month or year, picked at random, at each of seven United States stations are found to be estimated as closely, on the average, by the mean of the preceding 10 to 40 years as by a 30-year "normal." The median value for the preceding 15 or so years may be an even slightly better estimator. Graphs show the mean square and mean absolute differences between k -year means and the next observation for $k = 1$ [1] 50 , and the mean absolute differences between k -year medians and the next observation for $k = 1$ [2] 49 , for Dodge City, Vicksburg, Memphis, Cairo, Madison, Pittsburgh, and Lynchburg. Comparison of these graphs with corresponding graphs based on random normal numbers, biased in various ways, suggests that many climatic records contain progressive changes in mean or in variance, or both. The number of antecedent years (k^*) for which the mean or median is closest to the next year's observation varies erratically from month to month, but tends to be the same at nearby stations for a given month.

CLIMATIC NORMALS AS PREDICTORS, 2 : EXTENSION

1. Introduction

Confirmation, extension, and interpretation of the results of previous studies on the optimum averaging period for climatic data, when the average will be used to estimate next year's conditions, is the chief purpose of this Report. Further extensions and interpretations will be presented in future reports.

Scientific Report No. 1, henceforth designated as "SR 1", summarized and consolidated the findings of five previous studies. All were concerned with the average difference between a k -year average and a single value m years after the end of the k years, or an λ -year average beginning m years later:

$$i = \begin{array}{ccccccc} & & & \frac{k}{\dots\dots\dots} & \frac{m}{\text{---}} & \frac{\lambda}{\dots\dots\dots} & \\ & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & 2 & 3 & & & & n \end{array}$$

While this is the general formulation considered in the present study, the previous investigations were for single values ($\lambda = 1$) one year ahead ($m = 1$), except for one study in which the squared differences for prediction 1, 5, and 9 years ahead were averaged. For n observations ordered in time, x_1, x_2, \dots, x_n , the mean squared difference between a k -year average and the next ($k + 1$ st) value is

$$s_k^2 = \frac{1}{n - k} \sum_{i=1}^{n-k} \left[\frac{1}{k} \sum_{j=0}^{k-1} x_{i+j} - x_{i+k} \right]^2$$

This extrapolation variance S_k^2 and its square root, S_k , the standard error of extrapolation, were used by all except one of the previous investigators, and is the chief criterion of the present study. Two other plausible measures of extrapolation error, however, are also used:

Q_k , the mean prediction error, obtained by taking the absolute value, rather than the square, of the differences;

D_k , the mean prediction error of the median, defined as the mean absolute difference between the median of the antecedent k years and the $k + 1$ st value.

In the previous studies, discussed in SR 1, and in the present extension, the chief interest is in k^* ("k-star"), the value of k for which S_k^2 is a minimum. It is generally less than 30 years, the officially-adopted length of record for a climatic "normal," in the previous studies of daily maximum, mean monthly, and annual temperatures, of monthly and annual precipitation, and of annual streamflow. To confirm and extend these findings, monthly and annual temperatures and precipitation amounts for seven United States stations were used to compute S_k^2 and evaluate k^* ; other computations, for prediction more than one year ahead ($m > 1$), will be discussed in a later report.

The seven stations were all those in Vol. I (North America) of World Weather Records (1960) for which at least 85 years of complete data were given; actually all except one had 90 or more years. The seven, all in eastern or central United States, and the first year of complete data, are:

Dodge City, Kansas	1874
Vicksburg, Mississippi	1871
Memphis, Tennessee	1871
Cairo, Illinois	1871
Madison, Wisconsin	1869
Pittsburgh, Pennsylvania	1870
Lynchburg, Virginia	1871

All tables and figures present the data in this order, generally from west to east and south to north, up the Mississippi Valley.

Many U.S. stations have longer, and possibly "better," records. But the basic investigation seeks to establish what portion of a climatic record offers the best estimate of future conditions, at a reasonably good weather station, not a perfect one. These seven stations apparently had been deemed to have records sufficiently homogeneous and accurate to warrant world-wide use, and hence were suitable for study. Other stations, both in the U.S. and abroad, have been investigated, and will be discussed in future reports.

2. Discussion

For each of the seven stations, the mean squared prediction error, S_k^2 , was computed separately for monthly and annual temperature and precipitation, for $k = 1, 2, 3, \dots, 50$. The FORTRAN IV program, which also computes $S_{k/m}^2$ for $m = 1$ [1] 10, as well as $Q_{k/m}$ and $D_{k/m}$ and various ratios, is given in detail in Report No. 3. Results of these additional computations will be discussed later in this Report, and in subsequent reports.

As examples of the computational results, values of S_k^2 for annual temperature and precipitation at each of the seven stations are given in Tables 1 and 2 respectively. An asterisk indicates the minimum values of S_k^2 for each station, and hence identifies the appropriate k^* . All values of S_k^2 for each month, and the year, for each of the seven stations were plotted on the same basic diagrams, six months (or all stations, for annual values) to a sheet.

The basic plotting sheet, used in SR 1, has lines of $S_k^2 = (1 + 1/k) s^2$, for various arbitrary values of s^2 . If all the elements in a climatic series be normally distributed, independently and identically, with means of 0 and common variance s^2 , the difference between one observation and a

Table 1. Values of S_k^2 of annual temperatures at seven stations.

K	DODGE C.	VICKSB.	MEMPHIS	CAIRO	MADISON	PITTSB.	LYNCHB.
01	13.912	16.330	13.742	13.912	14.151	14.276	13.444
02	6.123	8.913	6.902	6.776	7.325	7.110	6.866
03	4.467	6.620	5.641	5.073	5.192	5.158	4.838
04	3.629	5.638	4.799	4.128	3.983	4.192	3.768
05	3.214	5.170	4.411	3.775	3.461	3.920	3.530
06	2.269	4.374	3.901	3.052	2.992	3.302	2.764
07	1.897	3.905	3.335	2.590	2.697	2.917	2.510
08	1.542	3.411	2.862	2.252	2.499	2.634	2.358
09	1.503	3.456	2.627	1.993	2.247	2.368	2.174
10	1.289	3.381	2.104	1.659	1.905	1.966	1.552
11	1.190*	3.405	1.906	1.489	1.748	1.739	1.576
12	1.248	3.256	1.768	1.465	1.751	1.727	1.652
13	1.356	3.092	1.601	1.388	1.680	1.679	1.591
14	1.406	3.043	1.434	1.445	1.669	1.697	1.671
15	1.359	3.004	1.427	1.420	1.628	1.661	1.701
16	1.325	2.998	1.294	1.404	1.598	1.612	1.736
17	1.321	2.909	1.165	1.456	1.680	1.743	1.733
18	1.279	2.803	1.170	1.160	1.430	1.442	1.530
19	1.300	2.613	1.111	1.044	1.347	1.359	1.282
20	1.340	2.544	0.980	1.014*	1.400	1.371	1.173
21	1.421	2.612	0.987	1.172	1.390	1.519	1.287
22	1.313	2.674	0.954	1.320	1.519	1.664	1.411
23	1.416	2.814	0.807	1.401	1.498	1.734	1.524
24	1.437	2.758	0.839	1.454	1.486	1.788	1.456
25	1.474	2.852	0.921	1.413	1.496	1.710	1.403
26	1.535	2.733	0.979	1.455	1.597	1.728	1.517
27	1.608	2.675	0.957	1.528	1.597	1.770	1.661
28	1.675	2.629	0.967	1.453	1.508	1.699	1.744
29	1.784	2.659	1.010	1.432	1.558	1.695	1.567
30	1.902	2.851	1.123	1.433	1.615	1.737	1.561
31	1.865	2.869	1.107	1.322	1.555	1.676	1.526
32	1.741	2.769	1.070	1.447	1.551	1.773	1.484
33	1.707	2.795	1.085	1.522	1.557	1.799	1.520
34	1.869	2.450	1.074	1.517	1.423	1.704	1.589
35	1.794	1.878	1.097	1.541	1.443	1.689	1.592
36	1.789	1.789	1.060	1.429	1.232	1.480	1.359
37	1.810	1.783	1.180	1.632	1.306	1.623	1.461
38	1.703	1.323	1.242	1.493	1.280	1.487	1.303
39	1.646	1.397	1.100	1.535	1.273	1.460	1.160
40	1.497	1.495	1.219	1.490	1.057*	1.349	1.163
41	1.365	1.389	1.179	1.525	1.065	1.386	1.189
42	1.513	1.309*	1.112	1.588	1.132	1.539	1.254
43	1.435	1.509	1.129	1.524	1.100	1.504	1.256
44	1.316	1.373	1.109	1.494	1.182	1.497	1.358
45	1.413	1.507	1.198	1.628	1.365	1.657	1.489
46	1.510	1.677	1.300	1.610	1.117	1.553	1.407
47	1.604	1.928	1.019	1.264	1.073	1.223*	1.095
48	1.916	1.469	1.025	1.385	1.168	1.372	1.097
49	2.104	1.700	0.874	1.570	1.304	1.568	1.184
50	2.006	1.713	0.624*	1.529	1.235	1.526	1.020*

Table 2. Values of S_k^2 of annual precipitation at seven stations.

K	DODGE C.	VICKSB.	MEMPHIS	CAIRO	MADISON	PITTSB.	LYNCHB.
01	23.592	17.844	15.928	12.692	17.177	13.620	17.290
02	14.503	11.203	9.297	6.746	9.393	7.375	11.203
03	10.156	8.312	7.040	4.523	7.195	4.898	7.321
04	8.631	6.722	5.914	3.450	6.135	3.816	5.945
05	7.956	6.184	5.532	2.784	5.873	3.234	5.278
06	7.248	5.460	4.217	2.299	5.611	2.755	4.300
07	6.292	5.016	3.826	2.080	5.477	2.583	4.214
08	6.218	4.792	3.620	1.865	5.563	2.278	3.775
09	6.231	4.469	3.1445	1.584	5.540	2.206	3.669
10	6.230	4.267	3.320	1.476	5.355	1.992	3.524
11	6.211	3.930	3.064	1.214	5.184	1.827	3.367
12	6.158	3.586	2.863	0.995	4.904	1.769	3.150
13	6.179	3.400	2.712	0.837	4.172	1.808	3.004
14	6.191	3.291	2.637	0.800	4.048	1.849	3.056
15	5.965	3.082	2.507	0.812	3.684	1.821	3.003
16	6.025	3.230	2.444	0.727	3.301	1.405	3.089
17	5.723	3.114	2.256	0.758	3.092	1.318	2.976
18	5.886	2.966	2.315	0.722*	2.935	1.339	2.594
19	5.865	3.038	2.259	0.793	2.534	1.038	2.568
20	5.824	3.010	2.294	0.830	2.517	0.990	2.558
21	5.713	3.037	2.244	0.948	2.478	1.020	2.594
22	5.668	3.069	1.923	0.857	2.410	1.061	2.453
23	5.580	3.200	1.835	0.915	2.374	0.995	2.475
24	5.322	3.296	1.808	0.965	2.298	0.977	2.403
25	4.998	3.252	1.838	0.918	2.307	0.936	2.264
26	4.893	3.064	1.872	0.936	2.284	0.945	2.144
27	4.991	3.050	1.837	0.837	2.098	0.934	1.873
28	5.255	3.002	1.882	0.984	1.964	1.100	1.794
29	4.960	2.817	1.844	1.055	2.029	1.127	1.863
30	5.070	2.789	1.750	1.140	1.959	0.773	1.315
31	5.087	2.712	1.841	1.122	2.026	0.819	1.217
32	4.408	2.691	1.923	1.308	1.997	0.878	1.210
33	4.557	2.760	1.952	1.439	1.912	...	1.151
34	4.408	2.919	2.002	1.693	1.960	1.038	1.046
35	3.697	3.031	1.776	1.604	1.837	1.123	1.059
36	3.820	2.955	1.843	1.744	1.783	1.080	0.751*
37	3.320	3.038	1.977	1.753	1.797	1.080	0.895
38	3.077	2.988	2.061	1.835	1.799	1.070	1.018
39	2.689	3.071	2.039	1.418	1.936	0.954	0.986
40	2.828	2.786	1.912	1.486	1.927	0.694*	0.981
41	2.681	2.745	2.027	1.667	1.513	0.757	0.909
42	2.768	2.612	2.095	1.584	1.245	0.845	1.012
43	2.876	2.536	2.074	1.744	1.158	0.917	1.041
44	1.365*	2.725	1.769	1.637	1.239	0.999	1.109
45	1.474	2.703	1.848	1.619	1.157	1.241	0.936
46	1.560	2.815	1.920	1.779	1.278	1.181	0.985
47	1.816	1.475	2.020	1.778	0.969*	1.327	0.975
48	1.597	1.357*	1.703	1.877	1.012	1.217	1.033
49	1.538	1.399	1.350	2.086	1.045	1.272	0.835
50	1.679	1.418	1.303*	2.056	1.230	1.331	0.931

k -element mean, not including it, is normally distributed with mean 0 and variance $s^2 + s^2/k$, as shown previously (SR 1). Hence S_k^2 should decrease as $(1 + 1/k) s^2$.

Departures from such behavior appear on all the figures. Minor fluctuations about the theoretical line could represent sampling errors: for two samples of 100 random normal numbers S_k^2 deviated somewhat from the appropriate line (SR 1, Figs. 8-11), but for 1,000 such numbers S_k^2 followed the line quite closely (SR 1, Fig. 13). Some of the large swings may be the results of changes in mean or in variance: S_k^2 increases with k if the mean changes, decreases if the variance changes (SR 1).

Thus a shift in mean is suggested for the annual temperatures (Fig. 8B) at all stations except possibly Vicksburg, as suggested for some stations in the actual data (Fig. 8A), graphed as departures from the mean. A shift in mean seems likely at Dodge City in February (Fig. 1A), and possibly at other stations in other months. In annual precipitation (Fig. 25B), Cairo shows great similarity to the most strongly biased synthetic case (SR 1, curve 4 of Figs. 8 and 10), with nearby Memphis somewhat less extreme; Lynchburg also has the same tendency. In the monthly precipitation curves, shifts in mean are suggested for:

Vicksburg, Fig. 19D, October and November

Memphis, Fig. 20C, January

Cairo, Fig. 21C, January

Madison, Fig. 22D, September

Lynchburg, Fig. 24D, August

Changes in variance, indicated by a plunging S_k^2 curve, are suggested by the curve for yearly precipitation at Madison (Fig. 24) and traces for:

Station	Temperature		Precipitation	
	Fig.	Month	Fig.	Month
Dodge City	1	Mar.	18	June., Sep.
Vicksburg			19	Oct., Dec.
Memphis	3	Jan., Dec.	20	Oct.
Cairo	4	Dec.	21	Oct.
Madison	5	Jan., Feb., Dec.	22	Jul.
Pittsburgh	6	Nov.	23	Apr.
Lynchburg			24	Sep., Oct.

Such shifts in mean or in variance, if actually the causes of the steep climbs or descents of the S_k^2 curves, do not come from changes in station location or observing practice. If they did, curves for consecutive months at the same station would be similar. But little coherence is shown by curves for successive months.

However, the S_k^2 curves for the same month at nearby stations often show marked similarities. Memphis and Cairo have already been noted as showing possible shifts in the January mean temperature, and variance shifts in December temperature and October precipitation. In fact, at these two stations, 150 miles apart along the Mississippi River, all corresponding S_k^2 curves for temperature are quite similar (Figs. 3 and 4), but the precipitation curves (Figs. 19 and 20) show no close similarities, although both have the same gross features in October.

This behavior agrees with the general climatologic principal that temperature has somewhat greater regional coherence than does precipitation. For each, the correlation at 150-mile separation is greater than the year-to-year correlation at any other place; for none of the seven stations is the serial correlation of either element in any month significantly different from zero, at any lag.

3. Comparisons

While the erratic behavior of S_k^2 , as k increases, poses interesting theoretical climatologic questions, it also raises an important practical problem. For each year at each of the seven stations, and for all of them together, what number (k) of antecedent years would have given a mean closest to that year's temperature and precipitation, for each month? A simple average of all the k^* values might not be appropriate, because many S_k^2 curves had such broad minima that $S_{k^*}^2$ was only fractionally less than values of S_k^2 for adjacent values of k .

In hopes of solving this problem, values of k^* and all other values of k for which S_k^2 was not more than 5 % greater than $S_{k^*}^2$ were graphed. All annual values were compared on rectangular diagrams, and monthly values at each place on polar diagrams, separately for temperature (Fig. 9) and for precipitation (Fig. 25). On each diagram, k^* is indicated by a dashed line, and single solid lines or shaded areas indicate those values of k for which $S_k^2 \leq 1.05 S_{k^*}^2$.

The only useful conclusion derived from many hours of study of these graphs, and of the S_k^2 curves, is that almost any value of k from 10 to 40 years would, on balance, give about the same predictive accuracy for these seven stations for the period studied. This tentative conclusion, or working hypothesis, will be examined, in later reports, with respect to other climatic elements and stations in other parts of the world, and for different periods.

First, however, comes the question of whether S_k^2 is an inherently unstable criterion. Perhaps the mean absolute error, Q_k , behaves better, and can more readily define the optimum period of record on which a climatic average or normal should be based, if it is to be used primarily for prediction

(which is really the primary use to which normals are put). Perhaps, also, a k-year median is a more suitable predictor, as measured by its mean absolute error, D_k , than is the k-year mean measured by either S_k^2 or Q_k .

Both Q_k and D_k were computed for annual and monthly temperature and precipitation at the same seven stations. Resulting values are shown in Figs. 10-17 and 27-34. These figures bear the same lines as the previous figures, showing the expected behavior of S_k^2 for independent numbers from a normal population. This is not the behavior expected of Q_k or D_k , which has as yet been established in theory. However, on computations with 1000 random normal variates, Slusser (SR 3, pp. 51-53) found that both Q_k and D_k reach their asymptotes much more quickly than does S_k^2 ; neither decreased appreciably after k exceeded 20 (Figs. 57 and 59, curves "O"). Hence in Figs. 10-17 and 27-34, the curves for Q_k and D_k would be expected to be somewhat flatter than the theoretical curves for S_k^2 , shown on the diagrams.

To eliminate ambiguity in defining the median, only odd values of k were used for D_k . Since Q_k , being also a mean absolute difference, is more suitable than S_k^2 for comparison with D_k , the graphs for Q_k were drawn for only the odd values of k , with the magnitude of Q_k for the even values of k indicated by dots, connected to the line.

No essential differences could be found between the behaviors of D_k , Q_k , and S_k^2 . In fact, for the same data the three curves are quite similar, rising, falling, jumping or steadying at the same values of k . The shifts in mean or in variance suggested by the S_k^2 behavior are not the consequences of taking squared differences from the mean, but are caused by inherent characteristics of the climatic data.

In almost two-thirds of the station-months, the smallest mean absolute difference from the median, D_{k^*} , was less than the smallest mean absolute difference from the mean, Q_{k^*} . In most cases, the number of years, k^* , was the same for the median as for the mean. But the median was somewhat closer, on the average, to the next year's value. Because of the similarity of behavior, no detailed analysis was made of the k^* values for Q_k or D_k . But the working hypothesis was expanded to suggest that the median of 10 to 40 antecedent years may be a better predictor than the mean for the same period, whose precise length seems immaterial. This hypothesis will be investigated further in subsequent reports.


4. Figures

All the properties of the monthly and annual temperature and precipitation at the seven U.S. stations are shown in 104 diagrams, grouped into 17 figures. Temperature diagrams are given first, followed by precipitation diagrams in exactly the same order. The entire arrangement is planned for easy intercomparisons.

In each series, a pair of monthly S_k^2 diagrams for each station is accompanied by a bar diagram of departures from normal. Annual values for all seven stations are shown as departures, and as S_k^2 curves. Behavior of k^* is then summarized on circular diagrams mounted on a map. Monthly graphs of Q_k and D_k for each station come next, followed by annual Q_k and D_k diagrams for all stations.

Figures exhibiting the behavior of temperature series at seven U.S. stations

Fig.	Page	Fig.	Page
1A, 1B, 1C	12	1B, 1C	13
2A, 2B, 2C	14	2B, 2C	15
3A, 3B, 3C	16	3B, 3C	17
4A, 4B, 4C	18	4B, 4C	19
5A, 5B, 5C	20	5B, 5C	21
6A, 6B, 6C	22	6B, 6C	23
7A, 7B, 7C	24	7B, 7C	25
8A, 8B	26	9	27
10A, 10B	28	10C, 10D	29
11A, 11B	30	11C, 11D	31
12A, 12B	32	12C, 12D	33
13A, 13B	34	13C, 13D	35
14A, 14B	36	14C, 14D	37
15A, 15B	38	15C, 15D	39
16A, 16B	40	16C, 16D,	41
17A, 17B	42		







FIGURE 18




FIGURE 2A





FIGURE 78






FIGURE 3A








FIGURE 44








FIGURE 54














FIGURE 6A






















FIGURE 11B




FIGURE 11A















FIGURE 11C











































Figures exhibiting the behavior of precipitation series at seven U.S. stations



Fig.		Page	Fig.		Page
18A, 18B	Precipitation departures, Dodge City	44	18C, 18D	S_k^2 of monthly precipitation, Dodge City	45
19A, 19B	" " Vicksburg	46	19C, 19D	" " Vicksburg	47
20A, 20B	" " Memphis	48	20C, 20D	" " Memphis	49
21A, 21B	" " Cairo	50	21C, 21D	" " Cairo	51
22A, 22B	" " Madison	52	22C, 22D	" " Madison	53
23A, 23B	" " Pittsburgh	54	23C, 23D	" " Pittsburgh	55
24A, 24B	" " Lynchburg	56	24C, 24D	" " Lynchburg	57
25A, 25B	Precipitation departures, annual. S_k^2 of annual precipitation, 7 stations.	58	26	k^* for S_k^2 of annual and monthly precipitation at seven stations.	59
27A, 27B	Q_k of monthly precipitation, Dodge City	60	27C, 27D	D_k of monthly precipitation, Dodge City	61
28A, 28B	" " Vicksburg	62	28C, 28D	" " Vicksburg	63
29A, 29B	" " Memphis	64	29C, 29D	" " Memphis	65
30A, 30B	" " Cairo	66	30C, 30D	" " Cairo	67
31A, 31B	" " Madison	68	31C, 31D	" " Madison	69
32A, 32B	" " Pittsburgh	70	32C, 32D	" " Pittsburgh	71
33A, 33B	" " Lynchburg	72	33C, 33D	" " Lynchburg	73
34A, 34B	Q_k and D_k of annual precipitation.	74			


















FIGURE 21B






FIGURE 21A






FIGURE 27B






FIGURE 27A






FIGURE 23B










FIGURE 23A










FIGURE 24B













FIGURE 24A























































UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author)		2a REPORT SECURITY CLASSIFICATION
SAN FERNANDO VALLEY STATE COLLEGE FOUNDATION Northridge, California 91324		UNCLASSIFIED
		2b GROUP
3 REPORT TITLE		
CLIMATIC NORMALS AS PREDICTORS. PART 2: EXTENSION		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates)		
Scientific. Interim.		
5 AUTHOR(S) (Last name, first name, initial)		
Arnold Court		
6 REPORT DATE	7a TOTAL NO. OF PAGES	7b NO. OF REFS
1 November 1968	76	none
8a CONTRACT OR GRANT NO.	8b. ORIGINATOR'S REPORT NUMBER(S)	
AF19(628)-5716	82-68-2	
b Project, Task, Work Unit Nos.	Scientific Report No. 2	
8624-TO2-WU 01		
c DoD Element 6240539F	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d DoD Subelement 681000	AFCRL-68-0673	
10 AVAILABILITY/LIMITATION NOTICES		
1-Distribution of this document is unlimited. It may be released to the Clearinghouse, Department of Commerce, for sale to the general public.		
11 SUPPLEMENTARY NOTES	12. SPONSORING MILITARY ACTIVITY	
TECH, OTHER	Air Force Cambridge Research Laboratories L. G. Hanscom Field (CRE) Bedford, Massachusetts 01730	
13 ABSTRACT		
<p>Temperature and precipitation for a single month or year, picked at random, at each of seven United States stations are found to be estimated as closely, on the average, by the mean of the preceding 10 to 40 years as by a 30-year "normal." The median value for the preceding 15 or so years may be an even slightly better estimator. Graphs show the mean square and mean absolute differences between k-year means and the next observation for $k = 1$ (1) 50, and the mean absolute differences between k-year medians and the next observation for $k = 1$ (2) 49, for Dodge City, Vicksburg, Memphis, Cairo, Madison, Pittsburgh, and Lynchburg. Comparison of these graphs with corresponding graphs based on random normal numbers, biased in various ways, suggests that many climatic records contain progressive changes in mean or in variance, or both. The number of antecedent years (k^*) for which the mean or median is closest to the next year's observation varies erratically from month to month, but tends to be the same at nearby stations for a given month.</p>		

DD FORM 1473
1 JAN 64

UNCLASSIFIED

Security Classification

UNCLASSIFIED
Security Classification

16. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
	NORMALS, CLIMATIC PREDICTION, CLIMATIC OPTIMUM RECORD LENGTH MEDIAN					

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, rules, and weights is optional.

UNCLASSIFIED
Security Classification